


Test Bank for

Maternal-Child Nursing 5th Edition

Authors: Emily McKinney, Susan James, Sharon Murray, Kristine Nelson, Jean Ashwill

- Chapter 1: Foundations of Maternity, Women's Health, and Child Health Nursing
- Chapter 2: The Nurse's Role in Maternity, Women's Health, and Pediatric Nursing
- Chapter 3: The Childbearing and Child-Rearing Family
- Chapter 4: Communicating with Children and Families
- Chapter 5: Health Promotion for the Developing Child
- Chapter 6: Health Promotion for the Infant
- Chapter 7: Health Promotion During Early Childhood
- Chapter 8: Health Promotion for the School-Age Child
- Chapter 9: Health Promotion for the Adolescent
- Chapter 10: Heredity and Environmental Influences on Development
- Chapter 11: Reproductive Anatomy and Physiology
- Chapter 12: Conception and Prenatal Development
- Chapter 13: Adaptations to Pregnancy
- Chapter 14: Nutrition for Childbearing
- Chapter 15: Prenatal Diagnostic Tests
- Chapter 16: Giving Birth
- Chapter 17: Intrapartum Fetal Surveillance
- Chapter 18: Pain Management for Childbirth
- Chapter 19: Nursing Care During Obstetric Procedures
- Chapter 20: Postpartum Adaptations
- Chapter 21: The Normal Newborn: Adaptation and Assessment
- Chapter 22: The Normal Newborn: Nursing Care
- Chapter 23: Newborn Feeding
- Chapter 24: The Childbearing Family with Special Needs
- Chapter 25: Pregnancy-Related Complications
- Chapter 26: Concurrent Disorders During Pregnancy
- Chapter 27: The Woman with an Intrapartum Complication
- Chapter 28: The Woman with a Postpartum Complication
- Chapter 29: The High-Risk Newborn: Problems Related to Gestational Age and Development
- Chapter 30: The High-Risk Newborn: Acquired and Congenital Conditions
- Chapter 31: Management of Fertility and Infertility
- Chapter 32: Women's Health Care
- Chapter 33: Physical Assessment of Children
- Chapter 34: Emergency Care of the Child
- Chapter 35: The Ill Child in the Hospital and Other Care Settings
- Chapter 36: The Child with a Chronic Condition or Terminal Illness
- Chapter 37: Principles and Procedures for Nursing Care of Children
- Chapter 38: Medication Administration and Safety for Infants and Children
- Chapter 39: Pain Management for Children
- Chapter 40: The Child with a Fluid and Electrolyte Alteration
- Chapter 41: The Child with an Infectious Disease
- Chapter 42: The Child with an Immunologic Alteration
- Chapter 43: The Child with a Gastrointestinal Alteration
- Chapter 44: The Child with a Genitourinary Alteration
- Chapter 45: The Child with a Respiratory Alteration
- Chapter 46: The Child with a Cardiovascular Alteration
- Chapter 47: The Child with a Hematologic Alteration
- Chapter 48: The Child with Cancer
- Chapter 49: The Child with an Alteration in Tissue Integrity
- Chapter 50: The Child with a Musculoskeletal Alteration
- Chapter 51: The Child with an Endocrine or Metabolic Alteration
- Chapter 52: The Child with a Neurologic Alteration
- Chapter 53: Psychosocial Problems in Children and Families



Chapter 01: Foundations of Maternity, Women's Health, and Child Health Nursing McKinney: Evolve Resources for Maternal-Child Nursing, 5th Edition

MULTIPLE CHOICE

1. Which factor significantly contributed to the shift from home births to hospital births in the early 20th century?
 - a. Puerperal sepsis was identified as a risk factor in labor and delivery.
 - b. Forceps were developed to facilitate difficult births.
 - c. The importance of early parental-infant contact was identified.
 - d. Technologic developments became available to physicians.

ANS: D

Technologic developments were available to physicians, not lay midwives. So in-hospital births increased in order to take advantage of these advancements. Puerperal sepsis has been a known problem for generations. In the late 19th century, Semmelweis discovered how it could be prevented with improved hygienic practices. The development of forceps is an example of a technology advance made in the early 20th century but is not the only reason birthplaces moved. Unlike home births, early hospital births hindered bonding between parents and their infants.

PTS: 1

DIF: Cognitive Level: Knowledge/Remembering

REF: p. 1

OBJ: Integrated Process: Teaching-Learning

MSC: Client Needs: Safe and Effective Care Environment

2. Family-centered maternity care developed in response to
 - a. demands by physicians for family involvement in childbirth.
 - b. the Sheppard-Towner Act of 1921.
 - c. parental requests that infants be allowed to remain with them rather than in a nursery.
 - d. changes in pharmacologic management of labor.

ANS: C

As research began to identify the benefits of early extended parent-infant contact, parents began to insist that the infant remain with them. This gradually developed into the practice of rooming-in and finally to family-centered maternity care. Family-centered care was a request by parents, not physicians. The Sheppard-Towner Act of 1921 provided funds for state-managed programs for mothers and children. The changes in pharmacologic management of labor were not a factor in family-centered maternity care.

PTS: 1

DIF: Cognitive Level: Knowledge/Remembering

REF: p. 2

OBJ: Integrated Process: Teaching-Learning

MSC: Client Needs: Psychosocial Integrity

3. Which setting for childbirth allows the least amount of parent-infant contact?
 - a. Labor/delivery/recovery/postpartum room
 - b. Birth center
 - c. Traditional hospital birth
 - d. Home birth

ANS: C

In the traditional hospital setting, the mother may see the infant for only short feeding periods, and the infant is cared for in a separate nursery. The labor/delivery/recovery/postpartum room setting allows increased parent-infant contact. Birth centers are set up to allow an increase in parent-infant contact. Home births allow an increase in parent-infant contact.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering
REF: p. 2 OBJ: Nursing Process: Planning
MSC: Client Needs: Health Promotion and Maintenance

4. As a result of changes in health care delivery and funding, a current trend seen in the pediatric setting is
- increased hospitalization of children.
 - decreased number of children living in poverty.
 - an increase in ambulatory care.
 - decreased use of managed care.

ANS: C

One effect of managed care has been that pediatric health care delivery has shifted dramatically from the acute care setting to the ambulatory setting in order to provide more cost-efficient care. The number of hospital beds being used has decreased as more care is given in outpatient settings and in the home. The number of children living in poverty has increased over the past decade. One of the biggest changes in health care has been the growth of managed care.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering
REF: p. 5 OBJ: Nursing Process: Planning
MSC: Client Needs: Safe and Effective Care Environment

5. The Women, Infants, and Children (WIC) program provides
- well-child examinations for infants and children living at the poverty level.
 - immunizations for high-risk infants and children.
 - screening for infants with developmental disorders.
 - supplemental food supplies to low-income pregnant or breastfeeding women.

ANS: D

WIC is a federal program that provides supplemental food supplies to low-income women who are pregnant or breastfeeding and to their children until age 5 years. Medicaid's Early and Periodic Screening, Diagnosis, and Treatment Program provides for well-child examinations and for treatment of any medical problems diagnosed during such checkups. Children in the WIC program are often referred for immunizations, but that is not the primary focus of the program. Public Law 99-457 is part of the Individuals with Disabilities Education Act that provides financial incentives to states to establish comprehensive early intervention services for infants and toddlers with, or at risk for, developmental disabilities.

PTS: 1 DIF: Cognitive Level: Comprehension REF: p. 8
OBJ: Integrated Process: Teaching-Learning
MSC: Client Needs: Health Promotion and Maintenance

6. In most states, adolescents who are not emancipated minors must have the permission of their parents before

- a. treatment for drug abuse.
- b. treatment for sexually transmitted diseases (STDs).
- c. accessing birth control.
- d. surgery.

ANS: D

Minors are not considered capable of giving informed consent, so a surgical procedure would require consent of the parent or guardian. Exceptions exist for obtaining treatment for drug abuse or STDs or for getting birth control in most states.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering
REF: p. 17 OBJ: Nursing Process: Planning
MSC: Client Needs: Safe and Effective Care Environment

7. The maternity nurse should have a clear understanding of the correct use of a clinical pathway. One characteristic of clinical pathways is that they
- a. are developed and implemented by nurses.
 - b. are used primarily in the pediatric setting.
 - c. set specific time lines for sequencing interventions.
 - d. are part of the nursing process.

ANS: C

Clinical pathways are standardized, interdisciplinary plans of care devised for patients with a particular health problem. They are used to identify patient outcomes, specify time lines to achieve those outcomes, direct appropriate interventions and sequencing of interventions, include interventions from a variety of disciplines, promote collaboration, and involve a comprehensive approach to care. They are developed by multiple health care professionals and reflect interdisciplinary care. They can be used in multiple settings and for patients throughout the life span. They are not part of the nursing process but can be used in conjunction with the nursing process to provide care to patients.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering
REF: p. 7 OBJ: Nursing Process: Planning
MSC: Client Needs: Safe and Effective Care Environment

8. The fastest growing group of homeless people is
- a. men and women preparing for retirement.
 - b. migrant workers.
 - c. single women and their children.
 - d. intravenous (IV) substance abusers.

ANS: C

Pregnancy and birth, especially for a teenager, are important contributing factors for becoming homeless. People preparing for retirement, migrant workers, and IV substance abusers are not among the fastest growing groups of homeless people.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering
REF: p. 14 OBJ: Nursing Process: Assessment
MSC: Client Needs: Physiologic Integrity

9. A nurse wishes to work to reduce infant mortality in the United States. Which activity would this nurse most likely participate in?
- Creating pamphlets in several different languages using an interpreter.
 - Assisting women to enroll in Medicaid by their third trimester.
 - Volunteering to provide prenatal care at community centers.
 - Working as an intake counselor at a women's shelter.

ANS: C

Prenatal care is vital to reducing infant mortality and medical costs. This nurse would most likely participate in community service providing prenatal care outreach activities in community centers, particularly in low-income areas. Pamphlets in other languages, enrolling in Medicaid, and working at a women's shelter all might impact infant mortality, but the greatest effect would be from assisting women to get consistent prenatal care.

PTS: 1 DIF: Cognitive Level: Application/Applying

REF: p. 14 OBJ: Nursing Process: Implementation

MSC: Client Needs: Health Promotion and Maintenance

10. The intrapartum woman sees no need for a routine admission fetal monitoring strip. If she continues to refuse, what is the first action the nurse should take?
- Consult the family of the woman.
 - Notify the provider of the situation.
 - Document the woman's refusal in the nurse's notes.
 - Make a referral to the hospital ethics committee.

ANS: B

Patients must be allowed to make choices voluntarily without undue influence or coercion from others. The physician, especially if unaware of the patient's decision, should be notified immediately. Both professionals can work to ensure the mother understands the rationale for the action and the possible consequences of refusal. The woman herself is the decision maker, unless incapacitated. Documentation should occur but is not the first action. This situation does not rise to the level of an ethical issue so there is no reason to call the ethics committee.

PTS: 1 DIF: Cognitive Level: Application/Applying

REF: p. 18 OBJ: Nursing Process: Implementation

MSC: Client Needs: Safe and Effective Care Environment

11. Which statement is true regarding the "quality assurance" or "incident" report?
- The report assures the legal department that no problem exists.
 - Reports are a permanent part of the patient's chart.
 - The nurse's notes should contain, "Incident report filed, and copy placed in chart."
 - This report is a form of documentation of an event that may result in legal action.

ANS: D

An incident report is used when something occurs that might result in legal action, such as a patient fall or medication error. It warns the legal department that there may be a problem in a particular patient's care. Incident reports are not part of the patient's chart; thus the nurses' notes should not contain any reference to them.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering

REF: p. 18 OBJ: Integrated Process: Communication and Documentation

MSC: Client Needs: Safe and Effective Care Environment

12. Elective abortion is considered an ethical issue because
- abortion law is unclear about a woman's constitutional rights.
 - the Supreme Court ruled that life begins at conception.
 - a conflict exists between the rights of the woman and the rights of the fetus.
 - it requires third-party consent.

ANS: C

Elective abortion is an ethical dilemma because two opposing courses of action are available. The belief that induced abortion is a private choice is in conflict with the belief that elective pregnancy termination is taking a life. Abortion laws are clear concerning a woman's constitutional rights. The Supreme Court has not ruled on when life begins. Abortion does not require third-party consent.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering

REF: p. 11 OBJ: Integrated Process: Teaching-Learning

MSC: Client Needs: Safe and Effective Care Environment

13. Which woman would be most likely to seek prenatal care?
- A 15-year-old who tells her friends, "I don't believe I'm pregnant."
 - A 20-year-old who is in her first pregnancy and has access to a free prenatal clinic.
 - A 28-year-old who is in her second pregnancy and abuses drugs and alcohol.
 - A 30-year-old who is in her fifth pregnancy and delivered her last infant at home.

ANS: B

The patient who acknowledges the pregnancy early, has access to health care, and has no reason to avoid health care is most likely to seek prenatal care. Being in denial about the pregnancy increases the risk of not seeking care. This patient is also 15, and other social factors may discourage her from seeking care as well. Women who abuse substances are less likely to receive prenatal care. Some women see pregnancy and delivery as a natural occurrence and do not seek health care.

PTS: 1 DIF: Cognitive Level: Comprehension/Understanding

REF: p. 14 OBJ: Nursing Process: Assessment

MSC: Client Needs: Health Promotion and Maintenance

14. A woman who delivered her baby 6 hours ago complains of headache and dizziness. The nurse administers an analgesic but does not perform any assessments. The woman then has a tonic-clonic seizure, falls out of bed, and fractures her femur. How would the actions of the nurse be interpreted in relation to standards of care?
- Negligent: the nurse failed to assess the woman for possible complications
 - Negligent: because the nurse medicated the woman
 - Not negligent: the woman had signed a waiver concerning the use of side rails
 - Not negligent: the woman did not inform the nurse of her symptoms as soon as they occurred

ANS: A

There are four elements to malpractice, which is negligence in the performance of professional duties: duty, breach of duty, damage, and proximate cause. The nurse was negligent because she or he did not perform any assessments, which is the first step of the nursing process and is a standard of care. By not assessing the patient, the nurse did not meet established standards of care, and thus is guilty of professional negligence, or malpractice.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering
REF: p. 16 OBJ: Nursing Process: Evaluation
MSC: Client Needs: Safe and Effective Care Environment

15. Which patient situation fails to meet the first requirement of informed consent?
- The patient does not understand the physician's explanations.
 - The physician gives the patient only a partial list of possible side effects and complications.
 - The patient is confused and disoriented.
 - The patient signs a consent form because her husband tells her to.

ANS: C

The first requirement of informed consent is that the patient must be competent to make decisions about health care. Full disclosure of information is an important element of the consent, but first the patient has to be competent to sign. Understanding is an important element of the consent, but first the patient has to be competent to sign. Voluntary consent is an important element of the consent, but first the patient has to be competent to sign.

PTS: 1 DIF: Cognitive Level: Knowledge/Remembering
REF: p. 17 OBJ: Nursing Process: Assessment
MSC: Client Needs: Safe and Effective Care Environment

16. Which situation reflects a potential ethical dilemma for the nurse?
- A nurse administers analgesics to a patient with cancer as often as the provider's order allows.
 - A neonatal nurse provides nourishment and care to a newborn who has a defect that is incompatible with life.
 - A labor nurse, whose religion opposes abortion, is asked to assist with an elective abortion.
 - A postpartum nurse provides information about adoption to a new mother who feels she cannot adequately care for her infant.

ANS: C

A dilemma exists in this situation because the nurse is being asked to assist with a procedure that she or he believes is morally wrong. The other situations do not contain elements of conflict for the nurse.

PTS: 1 DIF: Cognitive Level: Comprehension/Understanding
REF: p. 11 OBJ: Nursing Process: Assessment
MSC: Client Needs: Safe and Effective Care Environment

17. When planning a parenting class, the nurse should explain that the leading cause of death in children 1 to 4 years of age in the United States is
- premature birth.
 - congenital anomalies.