

# Test Bank Lehne's Pharmacology for Nursing Care, 11th Edition by Jacqueline Burchum, Laura Rosenthal

## Chapter 1: Orientation to Pharmacology Test Bank

---

### MULTIPLE CHOICE

1. The nurse is teaching a patient how a medication works to treat an illness. To do this, the nurse will rely on knowledge of:
  - a. clinical pharmacology.
  - b. drug efficacy.
  - c. pharmacokinetics.
  - d. pharmacotherapeutics.

ANS: D

Pharmacotherapeutics is the study of the use of drugs to diagnose, treat, and prevent conditions. Clinical pharmacology is concerned with all aspects of drug–human interactions. Drug efficacy measures the extent to which a given drug causes an intended effect. Pharmacokinetics is the study of the impact of the body on a drug.

DIF: Cognitive Level: Comprehension REF: Four Basic Terms

TOP: Nursing Process: Implementation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

2. What does it mean when a drug is described as easy to administer?
  - a. It can be stored indefinitely without need for refrigeration.
  - b. It does not interact significantly with other medications.
  - c. It enhances patient adherence to the drug regimen.
  - d. It is usually relatively inexpensive to produce.

ANS: C

A major benefit of drugs that are easy to administer is that patients taking them are more likely to comply with the drug regimen. Drugs that are easy to give may have the other attributes listed, but those properties are independent of ease of administration.

DIF: Cognitive Level: Comprehension

REF: Additional Properties of an Ideal Drug: Ease of Administration

TOP: Nursing Process: Assessment

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

3. A patient tells the nurse that he was told by the prescriber that the analgesic he is taking is very effective. Which statement by the patient demonstrates an understanding of the drug's effectiveness?
  - a. "I don't have to worry about toxicity, since it takes a large amount of this drug to cause an overdose."
  - b. "It has no side effects and doesn't interact with other drugs."
  - c. "I only have to take it every 12 hours."
  - d. "It might make me sleepy, and it lessens pain for several hours at a time."

ANS: D

A drug is effective if it produces the intended effects, even if it also produces side effects. Because no drug is completely safe, the level of toxicity does not determine effectiveness. All drugs have side effects and many react with other substances; these do not affect the drug's effectiveness. Ease of administration is independent of a drug's effectiveness.

DIF: Cognitive Level: Comprehension REF: Properties of an Ideal Drug

TOP: Nursing Process: Evaluation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

## MULTIPLE RESPONSE

1. What are the properties of an ideal drug? (Select all that apply.)
  - a. Irreversible action
  - b. Predictability
  - c. Ease of administration
  - d. Chemical stability
  - e. A simple trade name

ANS: B, C, D

In addition to predictability, ease of administration, and chemical stability, other properties include a reversible action so that any harm the drug may cause can be undone and a simple generic name, because generic names are usually complex and difficult to remember and pronounce.

DIF: Cognitive Level: Comprehension

REF: Properties of an Ideal Drug | Additional Properties of an Ideal Drug

TOP: Nursing Process: Assessment

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

2. Before administering a medication, what does the nurse need to know to evaluate how individual patient variability might affect the patient's response to the medication? (Select all that apply.)
  - a. Chemical stability of the medication
  - b. Ease of administration
  - c. Family medical history
  - d. Patient's age
  - e. Patient's diagnosis

ANS: C, D, E

The family medical history can indicate genetic factors that may affect a patient's response to a medication. Patients of different ages can respond differently to medications. The patient's illness can affect how drugs are metabolized. The chemical stability of the medication and the ease of administration are properties of drugs.

DIF: Cognitive Level: Analysis REF: Sources of Individual Variation

TOP: Nursing Process: Implementation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

## Chapter 2: Application of Pharmacology in Nursing Practice

### Test Bank

---

#### MULTIPLE CHOICE

1. A patient is using a metered-dose inhaler containing albuterol for asthma. The medication label instructs the patient to administer “2 puffs every 4 hours as needed for coughing or wheezing.” The patient reports feeling jittery sometimes when taking the medication, and she doesn’t feel that the medication is always effective. Which is *not* an appropriate nursing intervention for this patient?
  - a. Asking the patient to demonstrate use of the inhaler
  - b. Assessing the patient’s exposure to tobacco smoke
  - c. Auscultating lung sounds and obtaining vital signs
  - d. Suggesting that the patient use one puff to reduce side effects

ANS: D

It is not within the nurse’s scope of practice to change the dose of a medication without an order from a prescriber. Asking the patient to demonstrate inhaler use helps the nurse to evaluate the patient’s ability to administer the medication properly and is part of the nurse’s evaluation. Assessing tobacco smoke exposure helps the nurse determine whether nondrug therapies, such as smoke avoidance, can be used as an adjunct to drug therapy. Performing a physical assessment helps the nurse evaluate the patient’s response to the medication.

DIF: Cognitive Level: Application

REF: Applying the Nursing Process in Drug Therapy: Preadministration Assessment [and all subsections under this heading]

TOP: Nursing Process: Implementation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

2. A postoperative patient is being discharged home with acetaminophen/hydrocodone (Lortab) for pain. The patient asks the nurse about using Tylenol for fever. Which statement by the nurse is correct?
  - a. “It is not safe to take over-the-counter drugs with prescription medications.”
  - b. “Taking the two medications together poses a risk of drug toxicity.”
  - c. “There are no known drug interactions, so this will be safe.”
  - d. “Tylenol and Lortab are different drugs, so there is no risk of overdose.”

ANS: B

Tylenol is the trade name and acetaminophen is the generic name for the same medication. It is important to teach patients to be aware of the different names for the same drug to minimize the risk of overdose. Over-the-counter (OTC) medications and prescription medications may be taken together unless significant harmful drug interactions are possible. Even though no drug interactions are at play in this case, both drugs contain acetaminophen, which could lead to toxicity.

DIF: Cognitive Level: Application

REF: Application of Pharmacology in Patient Education: Dosage and Administration

TOP: Nursing Process: Implementation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

3. The nurse is preparing to care for a patient who will be taking an antihypertensive medication. Which action by the nurse is part of the assessment step of the nursing process?
- Asking the prescriber for an order to monitor serum drug levels
  - Monitoring the patient for drug interactions after giving the medication
  - Questioning the patient about over-the-counter medications
  - Taking the patient's blood pressure throughout the course of treatment

ANS: C

The assessment part of the nursing process involves gathering information before beginning treatment, and this includes asking about other medications the patient may be taking. Monitoring serum drug levels, watching for drug interactions, and checking vital signs after giving the medication are all part of the evaluation phase.

DIF: Cognitive Level: Application

REF: Preadministration Assessment

TOP: Nursing Process: Assessment

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

4. A postoperative patient reports pain, which the patient rates as an 8 on a scale from 1 to 10 (10 being the most extreme pain). The prescriber has ordered acetaminophen (Tylenol) 650 mg PO every 6 hours PRN pain. What will the nurse do?
- Ask the patient what medications have helped with pain in the past.
  - Contact the provider to request a different analgesic medication.
  - Give the pain medication and reposition the patient to promote comfort.
  - Request an order to administer the medication every 4 hours.

ANS: B

The nursing diagnosis for this patient is severe pain. Acetaminophen is given for mild to moderate pain, so the nurse should ask the prescriber to order a stronger analgesic medication. Asking the patient to tell the nurse what has helped in the past is a part of an initial assessment and should be done preoperatively and not when the patient is having severe pain. Because the patient is having severe pain, acetaminophen combined with nondrug therapies will not be sufficient. Increasing the frequency of the dose of a medication for mild pain will not be effective.

DIF: Cognitive Level: Analysis

REF: Analysis and Nursing Diagnosis

TOP: Nursing Process: Diagnosis

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

5. A patient newly diagnosed with diabetes is to be discharged from the hospital. The nurse teaching this patient about home management should begin by doing what?
- Asking the patient to demonstrate how to measure and administer insulin
  - Discussing methods of storing insulin and discarding syringes
  - Giving information about how diet and exercise affect insulin requirements
  - Teaching the patient about the long-term consequences of poor diabetes control

ANS: A

Because insulin must be given correctly to control symptoms and because an overdose can be fatal, it is most important for the patient to know how to administer it. Asking for a demonstration of technique is the best way to determine whether the patient has understood the teaching. When a patient is receiving a lot of new information, the information presented first is the most likely to be remembered. The other teaching points are important as well, but they are not as critical and can be taught later.

DIF: Cognitive Level: Application      REF: Planning      TOP: Nursing Process: Planning  
MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

6. The nurse receives an order to give morphine 5 mg IV every 2 hours PRN pain. Which action is *not* part of the six rights of drug administration?
- Assessing the patient's pain level 15 to 30 minutes after giving the medication
  - Checking the medication administration record to see when the last dose was administered
  - Consulting a drug manual to determine whether the amount the prescriber ordered is appropriate
  - Documenting the reason the medication was given in the patient's electronic medical record

ANS: A

Assessing the patient's pain after administering the medication is an important part of the nursing process when giving medications, but it is not part of the six rights of drug administration. Checking to see when the last dose was given helps ensure that the medication is given at the right time. Consulting a drug manual helps ensure that the medication is given in the right dose. Documenting the reason for a pain medication is an important part of the right documentation—the sixth right.

DIF: Cognitive Level: Application  
REF: Evolution of Nursing Responsibilities Regarding Drugs | Implementation  
TOP: Nursing Process: Implementation  
MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

7. A patient tells a nurse that a medication prescribed for recurrent migraine headaches is not working. What will the nurse do?
- Ask the patient about the number and frequency of tablets taken.
  - Assess the patient's headache pain on a scale from 1 to 10.
  - Report the patient's complaint to the prescriber.
  - Suggest biofeedback as an adjunct to drug therapy.

ANS: A

When evaluating the effectiveness of a drug, it is important to determine whether the patient is using the drug as ordered. Asking the patient to tell the nurse how many tablets are taken and how often helps the nurse determine compliance. Assessing current pain does not yield information about how well the medication is working unless the patient is currently taking it. The nurse should gather as much information about compliance, symptoms, and drug effectiveness as possible before contacting the prescriber. Biofeedback may be an effective adjunct to treatment, but it should not be recommended without complete information about drug effectiveness.

DIF: Cognitive Level: Application      REF: Evaluation      TOP: Nursing Process: Evaluation

8. A nurse is preparing to administer medications. Which patient would the nurse consider to have the greatest predisposition to an adverse reaction?
- A 30-year-old man with kidney disease
  - A 75-year-old woman with cystitis
  - A 50-year-old man with an upper respiratory tract infection
  - A 9-year-old boy with an ear infection

ANS: A

The individual with impaired kidney function would be at risk of having the drug accumulate to a toxic level because of potential excretion difficulties. Cystitis is an infection of the bladder and not usually the cause of excretion problems that might lead to an adverse reaction from a medication. A respiratory tract infection would not predispose a patient to an adverse reaction, because drugs are not metabolized or excreted by the lungs. A 9-year-old boy would not have the greatest predisposition to an adverse reaction simply because he is a child; nor does an ear infection put him at greater risk.

DIF: Cognitive Level: Analysis

REF: Minimizing Adverse Reactions

TOP: Nursing Process: Planning

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

9. A nurse consults a drug manual before giving a medication to an 80-year-old patient. The manual states that elderly patients are at increased risk for hepatic side effects. Which action by the nurse is correct?
- Contact the provider to discuss an order for pretreatment laboratory work.
  - Ensure that the drug is given in the correct dose at the correct time to minimize the risk of adverse effects.
  - Notify the provider that this drug is contraindicated for this patient.
  - Request an order to give the medication intravenously so that the drug does not pass through the liver.

ANS: A

The drug manual indicates that this drug should be given with caution to elderly patients. Getting information about liver function before giving the drug establishes baseline data that can be compared with post-treatment data to determine whether the drug is affecting the liver. Giving the correct dose at the correct interval helps to minimize risk, but without baseline information, the effects cannot be determined. The drug is not contraindicated.

DIF: Cognitive Level: Analysis

REF: Minimizing Adverse Effects

TOP: Nursing Process: Implementation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

10. A patient has been receiving intravenous penicillin for pneumonia for several days and begins to complain of generalized itching. The nurse auscultates bilateral wheezing and notes a temperature of 38.5° C (101° F). Which is the correct action by the nurse?
- Administer the next dose and continue to evaluate the patient's symptoms.
  - Ask the prescriber if an antihistamine can be given to relieve the itching.
  - Contact the prescriber to request an order for a chest radiograph.
  - Hold the next dose and notify the prescriber of the symptoms.

ANS: D

Pruritus and wheezing are signs of a possible allergic reaction, which can be fatal; therefore, the medication should not be given and the prescriber should be notified. When patients are having a potentially serious reaction to a medication, the nurse should not continue giving the medication. Antihistamines may help the symptoms of an allergic reaction, but the first priority is to stop the medication. Obtaining a chest radiograph is not helpful.

DIF: Cognitive Level: Application

REF: Evolution of Nursing Responsibilities Regarding Drugs | Application of Pharmacology in

Patient Care: Identifying High-Risk Patients

TOP: Nursing Process: Evaluation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

## **MULTIPLE RESPONSE**

1. A nurse is reviewing a patient's medical record before administering a medication. Which factor(s) can alter the patient's physiologic response to the drug? (Select all that apply.)
  - a. Ability to swallow pills
  - b. Age
  - c. Genetic factors
  - d. Gender
  - e. Height

ANS: B, C, D

Age, genetic factors, and gender all influence an individual patient's ability to absorb, metabolize, and excrete drugs; therefore, these factors must be assessed before a medication is administered. A patient's ability to swallow pills, although it may determine the way a drug is administered, does not affect the physiologic response. Height does not affect response; weight and the distribution of adipose tissue can affect the distribution of drugs.

DIF: Cognitive Level: Analysis

REF: Evolution of Nursing Responsibilities Regarding Drugs | Application of Pharmacology in

Patient Care: Preadministration Assessment

TOP: Nursing Process: Assessment

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

## Chapter 3: Drug Regulation, Development, Names, and Information

### Test Bank

---

#### MULTIPLE CHOICE

1. A nurse educator is conducting a continuing education class on pharmacology. To evaluate the learning of the nurses in the class, the nurse educator asks, "Which drug name is a generic drug name?" Which is the correct response?
  - a. Acetaminophen
  - b. Tylenol
  - c. Cipro
  - d. Motrin

ANS: A

Acetaminophen is the generic name. Tylenol, Cipro, and Motrin are all trade names.

DIF: Cognitive Level: Comprehension REF: Table 3-3: The Three Types of Drug Names

TOP: Nursing Process: Diagnosis

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

2. The FDA Amendments Act (FDAAA) was passed in 2007 to address which aspect of drug safety?
  - a. Allowing pharmaceutical companies to identify off-label uses of medications approved for other uses
  - b. Evaluating drug safety information that emerges after a drug has been approved and is in use
  - c. Expediting the approval process of the U.S. Food and Drug Administration (FDA) so that needed drugs can get to market more quickly
  - d. Requiring manufacturers to notify patients before removing a drug from the market

ANS: B

The FDAAA was passed to enable the Food and Drug Administration to continue oversight of a drug after granting it approval so that changes in labeling could be made as necessary and postmarketing risks could be tracked and identified. A provision of the FDA Modernization Act (FDAMA), passed in 1997, allows drug companies to promote their products for off-label uses as long as they promise to conduct studies to support their claims. Regulations to permit accelerated approval of drugs for life-threatening diseases were adopted in 1992 by the FDA. The requirement that drug companies notify patients 6 months before removing a drug from the market is a provision of the FDAMA.

DIF: Cognitive Level: Comprehension REF: Landmark Drug Legislation

TOP: Nursing Process: Evaluation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

3. A nursing student asks a nurse about pharmaceutical research and wants to know the purpose of randomization in drug trials. The nurse explains that randomization is used to:
  - a. ensure that differences in outcomes are the result of treatment and not differences in subjects.
  - b. compare the outcome caused by the treatment to the outcome caused by no treatment.

- c. make sure that researchers are unaware of which subjects are in which group.
- d. prevent subjects from knowing which group they are in and prevent preconception bias.

ANS: A

Randomization helps prevent allocation bias, which can occur when researchers place subjects with desired characteristics in the study group and other subjects in the control group so that differences in outcome are actually the result of differences in subjects and not treatment. Comparing treatment outcome to no treatment outcome is the definition of a controlled study. The last two options describe the use of blinding in studies; blinding ensures that researchers or subjects (or both) are unaware of which subjects are in which group so that preconceptions about benefits and risks cannot bias the results.

DIF: Cognitive Level: Comprehension REF: The Randomized Drug Trial

TOP: Nursing Process: Implementation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Pharmacologic and Parenteral Therapies

4. Someone asks a nurse about a new drug that is in preclinical testing and wants to know why it cannot be used to treat a friend's illness. Which statement by the nurse is correct?
- a. "A drug at this stage of development can be used only in patients with serious disease."
  - b. "At this stage of drug development, the safety and usefulness of the medication is unknown."
  - c. "Clinical trials must be completed to make sure the drug is safe to use in humans."
  - d. "Until postmarketing surveillance data are available, the drug cannot be used."

ANS: B

Preclinical testing must be completed before drugs can be tested in humans. In this stage, drugs are evaluated for toxicities, pharmacokinetic properties, and potentially useful effects. Some drugs can be used in patients before completion of Phase III studies, but this is after preclinical testing is complete. Clinical trials proceed in stages, and each stage has guidelines defining how a new drug may be used and which patients may receive it. Postmarketing surveillance takes place after a drug is in general use.

DIF: Cognitive Level: Comprehension

REF: Landmark Drug Legislation | Stages of Drug Development

TOP: Nursing Process: Implementation

MSC: NCLEX Client Needs Category: Physiologic Integrity: Reduction of Risk Potential

5. A patient asks a nurse why drugs that have been approved by the FDA still have unknown side effects. The nurse tell the patient that:
- a. testing for all side effects of a medication would be prohibitively expensive.
  - b. patients in drug trials often are biased by their preconceptions of a drug's benefits.
  - c. researchers tend to conduct studies that will prove the benefits of their new drugs.
  - d. subjects in drug trials do not always represent the full spectrum of possible patients.

ANS: D